

**JEE-2007**  
**Paper 2**

1. In the experiment to determine the speed of sound using a resonance column,
- (A) prongs of the tuning fork are kept in a vertical plane
  - (B) prongs of the tuning fork are kept in a horizontal plane
  - (C) in one of the two resonances observed, the length of the resonating air column is close to the wavelength of sound in air
  - (D) in one of the two resonances observed, the length of the resonating air column is close to half of the wavelength of sound in air

Answer

- ☒ ☐ ☐ ☐  
(A) (B) (C) (D)

2. A student performs an experiment to determine the Young's modulus of a wire, exactly 2 m long, by Searle's method. In a particular reading, the student measures the extension in the length of the wire to be 0.8 mm with an uncertainty of  $\pm 0.05$  mm at a load of exactly 1.0 kg. The student also measures the diameter of the wire to be 0.4 mm with an uncertainty of  $\pm 0.01$  mm. Take  $g = 9.8 \text{ m/s}^2$  (exact). The Young's modulus obtained from the reading is
- (A)  $(2.0 \pm 0.3) \times 10^{11} \text{ N/m}^2$
  - (B)  $(2.0 \pm 0.2) \times 10^{11} \text{ N/m}^2$
  - (C)  $(2.0 \pm 0.1) \times 10^{11} \text{ N/m}^2$
  - (D)  $(2.0 \pm 0.05) \times 10^{11} \text{ N/m}^2$

Answer

- ☒ ☐ ☐ ☐  
(A) (B) (C) (D)

**OR**


- ☐ ☒ ☐ ☐  
(A) (B) (C) (D)

3. A particle moves in the X-Y plane under the influence of a force such that its linear momentum is  $\vec{p}(t) = A [\hat{i} \cos(kt) - \hat{j} \sin(kt)]$ , where  $A$  and  $k$  are constants. The angle between the force and the momentum is
- (A)  $0^\circ$
  - (B)  $30^\circ$
  - (C)  $45^\circ$
  - (D)  $90^\circ$

Answer

- ☐ ☐ ☐ ☒  
(A) (B) (C) (D)

4. A small object of uniform density rolls up a curved surface with an initial velocity  $v$ . It reaches up to a maximum height of  $\frac{3v^2}{4g}$  with respect to the initial position. The object is




- (A) ring      (B) solid sphere      (C) hollow sphere      (D) disc

Answer

- ☐ (A)   
 ☐ (B)   
 ☐ (C)   
 ☒ (D)

5. Water is filled up to a height  $h$  in a beaker of radius  $R$  as shown in the figure. The density of water is  $\rho$ , the surface tension of water is  $T$  and the atmospheric pressure is  $P_0$ . Consider a vertical section ABCD of the water column through a diameter of the beaker. The force on water on one side of this section by water on the other side of this section has magnitude




- (A)  $|2P_0 R h + \pi R^2 \rho g h - 2RT|$       (B)  $|2P_0 R h + R \rho g h^2 - 2RT|$   
 (C)  $|P_0 \pi R^2 + R \rho g h^2 - 2RT|$       (D)  $|P_0 \pi R^2 + R \rho g h^2 + 2RT|$

Answer

- ☐ (A)   
 ☒ (B)   
 ☐ (C)   
 ☐ (D)

6. A spherical portion has been removed from a solid sphere having a charge distributed uniformly in its volume as shown in the figure. The electric field inside the emptied space is








- (A) zero everywhere                      (B) non-zero and uniform  
(C) non-uniform                      (D) zero only at its center

Answer ☐ ☒ ☐ ☐  
(A) (B) (C) (D)


7. Positive and negative point charges of equal magnitude are kept at  $\left(0, 0, \frac{a}{2}\right)$  and  $\left(0, 0, -\frac{a}{2}\right)$ , respectively. The work done by the electric field when another positive point charge is moved from  $(-a, 0, 0)$  to  $(0, a, 0)$  is
- (A) positive  
(B) negative  
(C) zero  
(D) depends on the path connecting the initial and final positions

Answer ☐ ☐ ☒ ☐  
(A) (B) (C) (D)

8. A magnetic field  $\vec{B} = B_0 \hat{j}$  exists in the region  $a < x < 2a$  and  $\vec{B} = -B_0 \hat{j}$ , in the region  $2a < x < 3a$ , where  $B_0$  is a positive constant. A positive point charge moving with a velocity  $\vec{v} = v_0 \hat{i}$ , where  $v_0$  is a positive constant, enters the magnetic field at  $x = a$ . The trajectory of the charge in this region can be like,



Answer



9. Electrons with de-Broglie wavelength  $\lambda$  fall on the target in an X-ray tube. The cut-off wavelength of the emitted X-rays is


(A)  $\lambda_0 = \frac{2mc\lambda^2}{h}$

(B)  $\lambda_0 = \frac{2h}{mc}$

(C)  $\lambda_0 = \frac{2m^2 c^2 \lambda^3}{h^2}$

(D)  $\lambda_0 = \lambda$

Answer





10. STATEMENT-1

If there is no external torque on a body about its center of mass, then the velocity of the center of mass remains constant.

**because**

STATEMENT-2

The linear momentum of an isolated system remains constant.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer



(A)



(B)



(C)



(D)

11. STATEMENT-1

A cloth covers a table. Some dishes are kept on it. The cloth can be pulled out without dislodging the dishes from the table.

**because**

STATEMENT-2

For every action there is an equal and opposite reaction.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer



(A)



(B)



(C)



(D)


12. STATEMENT-1

A vertical iron rod has a coil of wire wound over it at the bottom end. An alternating current flows in the coil. The rod goes through a conducting ring as shown in the figure. The ring can float at a certain height above the coil.

**because**

STATEMENT-2

In the above situation, a current is induced in the ring which interacts with the horizontal component of the magnetic field to produce an average force in the upward direction.



- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

**Answer**

- ☒ (A)
 ☐ (B)
 ☐ (C)
 ☐ (D)

13. STATEMENT-1

The total translational kinetic energy of all the molecules of a given mass of an ideal gas is 1.5 times the product of its pressure and its volume.

**because**

STATEMENT-2

The molecules of a gas collide with each other and the velocities of the molecules change due to the collision.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

**Answer**


- ☐ (A)
 ☒ (B)
 ☐ (C)
 ☐ (D)

14. The speed of sound of the whistle is
- (A) 340 m/s for passengers in A and 310 m/s for passengers in B
  - (B) 360 m/s for passengers in A and 310 m/s for passengers in B
  - (C) 310 m/s for passengers in A and 360 m/s for passengers in B
  - (D) 340 m/s for passengers in both the trains

Answer ☐ ☒ ☐ ☐

(A) (B) (C) (D)

15. The distribution of the sound intensity of the whistle as observed by the passengers in train A is best represented by



Answer ☒ ☐ ☐ ☐

(A) (B) (C) (D)

16. The spread of frequency as observed by the passengers in train B is
- (A) 310 Hz
  - (B) 330 Hz
  - (C) 350 Hz
  - (D) 290 Hz

Answer ☒ ☐ ☐ ☐

(A) (B) (C) (D)

17. Light travels as a
- (A) parallel beam in each medium
  - (B) convergent beam in each medium
  - (C) divergent beam in each medium
  - (D) divergent beam in one medium and convergent beam in the other medium

Answer ☒ ☐ ☐ ☐

(A) (B) (C) (D)

18. The phases of the light wave at  $c, d, e$  and  $f$  are  $\phi_c, \phi_d, \phi_e$  and  $\phi_f$  respectively.

It is given that  $\phi_c \neq \phi_f$ .

- (A)  $\phi_c$  cannot be equal to  $\phi_d$                       (B)  $\phi_d$  can be equal to  $\phi_e$   
(C)  $(\phi_d - \phi_f)$  is equal to  $(\phi_c - \phi_e)$                       (D)  $(\phi_d - \phi_c)$  is not equal to  $(\phi_f - \phi_e)$

Answer

- ☐ ☐ ☒ ☐  
(A)      (B)      (C)      (D)

19. Speed of light is

- (A) the same in medium-1 and medium-2  
(B) larger in medium-1 than in medium-2  
(C) larger in medium-2 than in medium-1  
(D) different at  $b$  and  $d$

Answer

- ☐ ☒ ☐ ☐  
(A)      (B)      (C)      (D)

20. **Column I** describes some situations in which a small object moves. **Column II** describes some characteristics of these motions. Match the situations in **Column I** with the characteristics in **Column II** and indicate your answer by darkening appropriate bubbles in the  $4 \times 4$  matrix given in the ORS.

**Column I**

**Column II**

- |   |  |
|---|--|
| <p>(A) The object moves on the <math>x</math>-axis under a conservative force in such a way that its “speed” and “position” satisfy <math>v = c_1 \sqrt{c_2 - x^2}</math>, where <math>c_1</math> and <math>c_2</math> are positive constants.</p> <p>(B) The object moves on the <math>x</math>-axis in such a way that its velocity and its displacement from the origin satisfy <math>v = -kx</math>, where <math>k</math> is a positive constant.</p> <p>(C) The object is attached to one end of a mass-less spring of a given spring constant. The other end of the spring is attached to the ceiling of an elevator. Initially everything is at rest. The elevator starts going upwards with a constant acceleration <math>a</math>. The motion of the object is observed from the elevator during the period it maintains this acceleration.</p> <p>(D) The object is projected from the earth’s surface vertically upwards with a speed <math>2\sqrt{GM_e/R_e}</math>, where, <math>M_e</math> is the mass of the earth and <math>R_e</math> is the radius of the earth. Neglect forces from objects other than the earth.</p> | <p>(p) The object executes a simple harmonic motion.</p> <p>(q) The object does not change its direction.</p> <p>(r) The kinetic energy of the object keeps on decreasing.</p> <p>(s) The object can change its direction only once.</p> |
|---|--|

**Answer**

|   | p                                | q                                | r                                | s                     |
|---|----------------------------------|----------------------------------|----------------------------------|-----------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/> |
| B | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |
| C | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/> |
| D | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |


**OR**

|   | p                                | q                                | r                                | s                     |
|---|----------------------------------|----------------------------------|----------------------------------|-----------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/> |
| B | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |
| C | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/> |
| D | <input type="radio"/>            | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/> |


21. Two wires each carrying a steady current  $I$  are shown in four configurations in **Column I**. Some of the resulting effects are described in **Column II**. Match the statements in **Column I** with the statements in **Column II** and indicate your answer by darkening appropriate bubbles in the  $4 \times 4$  matrix given in the ORS.

**Column I**


- (A) Point P is situated midway between the wires.




- (B) Point P is situated at the mid-point of the line joining the centers of the circular wires, which have same radii.



- (C) Point P is situated at the mid-point of the line joining the centers of the circular wires, which have same radii.



- (D) Point P is situated at the common center of the wires.



**Column II**

- (p) The magnetic fields ( $B$ ) at P due to the currents in the wires are in the same direction.

- (q) The magnetic fields ( $B$ ) at P due to the currents in the wires are in opposite directions.

- (r) There is no magnetic field at P.

- (s) The wires repel each other.

**Answer**

|   | p                                | q                                | r                                | s                     |
|---|----------------------------------|----------------------------------|----------------------------------|-----------------------|
| A | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |
| B | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/> |
| C | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |
| D | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/> |

**OR**

|   | p                                | q                                | r                                | s                                |
|---|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| A | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            |
| B | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/>            |
| C | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            |
| D | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            | <input checked="" type="radio"/> |

22. **Column I** gives some devices and **Column II** gives some processes on which the functioning of these devices depend. Match the devices in **Column I** with the processes in **Column II** and indicate your answer by darkening appropriate bubbles in the  $4 \times 4$  matrix given in the ORS.

**Column I**

- (A) Bimetallic strip  
(B) Steam engine  
(C) Incandescent lamp  
(D) Electric fuse

**Column II**

- (p) Radiation from a hot body  
(q) Energy conversion  
(r) Melting  
(s) Thermal expansion of solids

**Answer** A – ‘s, q’ OR ‘s’ alone

B – ‘q’

C – ‘p, q’ OR ‘p’ alone

D – ‘q, r’ OR ‘r’ alone

23. Consider a titration of potassium dichromate solution with acidified Mohr’s salt solution using diphenylamine as indicator. The number of moles of Mohr’s salt required per mole of dichromate is

- (A) 3 (B) 4 (C) 5 (D) 6

**Answer** ☐ ☐ ☐ ☒  
(A) (B) (C) (D)

24. Among the following metal carbonyls, the C–O bond order is lowest in

- (A)  $[\text{Mn}(\text{CO})_6]^+$  (B)  $[\text{Fe}(\text{CO})_5]$  (C)  $[\text{Cr}(\text{CO})_6]$  (D)  $[\text{V}(\text{CO})_6]^-$





**Answer** ☐ ☐ ☐ ☒  
(A) (B) (C) (D)

25. A solution of a metal ion when treated with KI gives a red precipitate which dissolves in excess KI to give a colourless solution. Moreover, the solution of metal ion on treatment with a solution of cobalt(II) thiocyanate gives rise to a deep blue crystalline precipitate. The metal ion is


- (A)  $\text{Pb}^{2+}$  (B)  $\text{Hg}^{2+}$  (C)  $\text{Cu}^{2+}$  (D)  $\text{Co}^{2+}$

**Answer** ☐ ☒ ☐ ☐  
(A) (B) (C) (D)

26. Cyclohexene on ozonolysis followed by reaction with zinc dust and water gives compound **E**. Compound **E** on further treatment with aqueous KOH yields compound **F**. Compound **F** is




Answer







27. The number of stereoisomers obtained by bromination of *trans*-2-butene is

(A) 1 (B) 2 (C) 3 (D) 4


Answer



28. Among the following, the least stable resonance structure is



Answer





29. A positron is emitted from  $^{23}_{11}\text{Na}$ . The ratio of the atomic mass and atomic number of the resulting nuclide is

(A) 22/10

(B) 22/11

(C) 23/10

(D) 23/12

Answer



(A)



(B)



(C)



(D)

30. For the process  $\text{H}_2\text{O}(l)$  (1 bar, 373 K)  $\rightarrow$   $\text{H}_2\text{O}(g)$  (1 bar, 373 K), the correct set of thermodynamic parameters is

(A)  $\Delta G = 0$ ,  $\Delta S = +ve$

(B)  $\Delta G = 0$ ,  $\Delta S = -ve$

(C)  $\Delta G = +ve$ ,  $\Delta S = 0$

(D)  $\Delta G = -ve$ ,  $\Delta S = +ve$

Answer



(A)



(B)



(C)



(D)

31. Consider a reaction  $aG + bH \rightarrow \text{Products}$ . When concentration of both the reactants G and H is doubled, the rate increases by eight times. However, when concentration of G is doubled keeping the concentration of H fixed, the rate is doubled. The overall order of the reaction is

(A) 0

(B) 1

(C) 2

(D) 3

Answer



(A)



(B)



(C)



(D)

32. STATEMENT-1 : Alkali metals dissolve in liquid ammonia to give blue solutions.

**because**

STATEMENT-2 : Alkali metals in liquid ammonia give solvated species of the type  $[M(NH_3)_n]^+$  (M = alkali metals).

- (A) Statement-1 is True, Statement-2 is True; Statement-2 **is** a correct explanation for Statement-1
- (B) Statement-1 is True; Statement-2 is True; Statement-2 **is NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

**Answer**

- ☐ ☒ ☐ ☐  
(A) (B) (C) (D)

33. STATEMENT-1 : Glucose gives a reddish-brown precipitate with Fehling's solution.

**because**

STATEMENT-2 : Reaction of glucose with Fehling's solution gives CuO and gluconic acid.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 **is** a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 **is NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

**Answer**

- ☐ ☐ ☒ ☐  
(A) (B) (C) (D)

34. STATEMENT-1 : Molecules that are not superimposable on their mirror images are chiral.

because

STATEMENT-2 : All chiral molecules have chiral centres.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer

- ☐ ☐ ☒ ☐
- (A) (B) (C) (D)

35. STATEMENT-1 : Band gap in germanium is small.

because

STATEMENT-2 : The energy spread of each germanium atomic energy level is infinitesimally small.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer

- ☐ ☒ ☐ ☐
- (A) (B) (C) (D)

36. Among the following, identify the correct statement.

- (A) Chloride ion is oxidised by  $O_2$       (B)  $Fe^{2+}$  is oxidised by iodine  
(C) Iodide ion is oxidised by chlorine      (D)  $Mn^{2+}$  is oxidised by chlorine

Answer

- ☐ (A)    ☐ (B)    ☒ (C)    ☐ (D)

37. While  $Fe^{3+}$  is stable,  $Mn^{3+}$  is not stable in acid solution because

- (A)  $O_2$  oxidises  $Mn^{2+}$  to  $Mn^{3+}$   
(B)  $O_2$  oxidises both  $Mn^{2+}$  to  $Mn^{3+}$  and  $Fe^{2+}$  to  $Fe^{3+}$   
(C)  $Fe^{3+}$  oxidises  $H_2O$  to  $O_2$   
(D)  $Mn^{3+}$  oxidises  $H_2O$  to  $O_2$

Answer

- ☐ (A)    ☐ (B)    ☐ (C)    ☒ (D)

38. Sodium fusion extract, obtained from aniline, on treatment with iron(II) sulphate and  $H_2SO_4$  in presence of air gives a Prussian blue precipitate. The blue colour is due to the formation of

- (A)  $Fe_4[Fe(CN)_6]_3$                       (B)  $Fe_3[Fe(CN)_6]_2$   
(C)  $Fe_4[Fe(CN)_6]_2$                       (D)  $Fe_3[Fe(CN)_6]_3$

Answer

- ☒ (A)    ☐ (B)    ☐ (C)    ☐ (D)

39. Which one of the following reagents is used in the above reaction?

- (A) aq.  $NaOH + CH_3Cl$                       (B) aq.  $NaOH + CH_2Cl_2$   
(C) aq.  $NaOH + CHCl_3$                       (D) aq.  $NaOH + CCl_4$

Answer

- ☐ (A)    ☐ (B)    ☒ (C)    ☐ (D)





10. The electrophile in this reaction is

- (A)  $\text{:CHCl}$       (B)  $^+\text{CHCl}_2$       (C)  $\text{:CCl}_2$       (D)  $\cdot\text{CCl}_3$

Answer

- ☐ (A)    ☐ (B)    ☒ (C)    ☐ (D)

41. The structure of the intermediate I is



Answer

- ☐ (A)    ☒ (B)    ☐ (C)    ☐ (D)

42. Match the reactions in **Column I** with nature of the reactions/type of the products in **Column II**. Indicate your answer by darkening the appropriate bubbles of the  $4 \times 4$  matrix given in the ORS.

**Column I**

**Column II**

- |  |   |
|--|---|
| (A) $\text{O}_2^- \rightarrow \text{O}_2 + \text{O}_2^{2-}$              | (p) redox reaction                                    |
| (B) $\text{CrO}_4^{2-} + \text{H}^+ \rightarrow$                         | (q) one of the products has trigonal planar structure |
| (C) $\text{MnO}_4^- + \text{NO}_2^- + \text{H}^+ \rightarrow$            | (r) dimeric bridged tetrahedral metal ion             |
| (D) $\text{NO}_3^- + \text{H}_2\text{SO}_4 + \text{Fe}^{2+} \rightarrow$ | (s) disproportionation                                |

**Answer**

|   | p                                | q                                | r                                | s                                |
|---|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input checked="" type="radio"/> |
| B | <input type="radio"/>            | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            |
| C | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            |
| D | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/>            |

43. Match the compounds/ions in **Column I** with their properties/reactions in **Column II**. Indicate your answer by darkening the appropriate bubbles of the  $4 \times 4$  matrix given in the ORS.

**Column I**

**Column II**

- |  |   |
|--|---|
| (A) $\text{C}_6\text{H}_5\text{CHO}$     | (p) gives precipitate with 2,4-dinitrophenylhydrazine |
| (B) $\text{CH}_3\text{C}\equiv\text{CH}$ | (q) gives precipitate with $\text{AgNO}_3$            |
| (C) $\text{CN}^-$                        | (r) is a nucleophile                                  |
| (D) $\text{I}^-$                         | (s) is involved in cyanohydrin formation              |

**Answer**

|   | p                                | q                                | r                                | s                                |
|---|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input checked="" type="radio"/> |
| B | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            |
| C | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input checked="" type="radio"/> |
| D | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            |

44. Match the crystal system/unit cells mentioned in **Column I** with their characteristic features mentioned in **Column II**. Indicate your answer by darkening the appropriate bubbles of the  $4 \times 4$  matrix given in the ORS.

| Column I                                | Column II  |
|---|--|
| (A) simple cubic and face-centred cubic | (p) have these cell parameters $a=b=c$ and $\alpha = \beta = \gamma$ |
| (B) cubic and rhombohedral              | (q) are two crystal systems  |
| (C) cubic and tetragonal                | (r) have only two crystallographic angles of $90^\circ$              |
| (D) hexagonal and monoclinic            | (s) belong to same crystal system                                    |

Answer

|   | p                                | q                                | r                                | s                                |
|---|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            | <input checked="" type="radio"/> |
| B | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            |
| C | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/>            |
| D | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            |

45. Let  $O(0, 0)$ ,  $P(3, 4)$ ,  $Q(6, 0)$  be the vertices of the triangle  $OPQ$ . The point  $R$  inside the triangle  $OPQ$  is such that the triangles  $OPR$ ,  $PQR$ ,  $OQR$  are of equal area. The coordinates of  $R$  are

- |                                   |   |
|-----------------------------------|---|
| (A) $\left(\frac{4}{3}, 3\right)$ | (B) $\left(3, \frac{2}{3}\right)$           |
| (C) $\left(3, \frac{4}{3}\right)$ | (D) $\left(\frac{4}{3}, \frac{2}{3}\right)$ |

Answer

- |                       |                       |                                  |                       |
|-----------------------|-----------------------|----------------------------------|-----------------------|
| <input type="radio"/> | <input type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |
| (A)                   | (B)                   | (C)                              | (D)                   |

46. If  $|z| = 1$  and  $z \neq \pm 1$ , then all the values of  $\frac{z}{1-z^2}$  lie on

- (A) a line not passing through the origin
- (B)  $|z| = \sqrt{2}$
- (C) the  $x$ -axis
- (D) the  $y$ -axis

Answer



(A) (B) (C) (D)

47. Let  $E^c$  denote the complement of an event  $E$ . Let  $E, F, G$  be pairwise independent events with  $P(G) > 0$  and  $P(E \cap F \cap G) = 0$ . Then  $P(E^c \cap F^c | G)$  equals

- (A)  $P(E^c) + P(F^c)$
- (B)  $P(E^c) - P(F^c)$
- (C)  $P(E^c) - P(F)$
- (D)  $P(E) - P(F^c)$

Answer



(A) (B) (C) (D)

48.  $\frac{d^2x}{dy^2}$  equals

- (A)  $\left(\frac{d^2y}{dx^2}\right)^{-1}$
- (B)  $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$
- (C)  $\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$
- (D)  $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

Answer




(A) (B) (C) (D)




49. The differential equation  $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{y}$  determines a family of circles with
- (A) variable radii and a fixed centre at  $(0, 1)$   
 (B) variable radii and a fixed centre at  $(0, -1)$   
 (C) fixed radius 1 and variable centres along the  $x$ -axis  
 (D) fixed radius 1 and variable centres along the  $y$ -axis

Answer




50. Let  $\vec{a}, \vec{b}, \vec{c}$  be unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ . Which one of the following is correct?
- (A)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} = \vec{0}$   
 (B)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \neq \vec{0}$   
 (C)  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{a} \times \vec{c} \neq \vec{0}$   
 (D)  $\vec{a} \times \vec{b}, \vec{b} \times \vec{c}, \vec{c} \times \vec{a}$  are mutually perpendicular

Answer




51. Let  $ABCD$  be a quadrilateral with area 18, with side  $AB$  parallel to the side  $CD$  and  $AB = 2CD$ . Let  $AD$  be perpendicular to  $AB$  and  $CD$ . If a circle is drawn inside the quadrilateral  $ABCD$  touching all the sides, then its radius is
- (A) 3                      (B) 2                      (C)  $\frac{3}{2}$                       (D) 1

Answer




52. Let  $f(x) = \frac{x}{(1+x^n)^{1/n}}$  for  $n \geq 2$  and  $g(x) = \underbrace{(f \circ f \circ \dots \circ f)}_{f \text{ occurs } n \text{ times}}(x)$ . Then  $\int x^{n-2} g(x) dx$  equals
- (A)  $\frac{1}{n(n-1)}(1+nx^n)^{1-\frac{1}{n}} + K$  (B)  $\frac{1}{n-1}(1+nx^n)^{1-\frac{1}{n}} + K$
- (C)  $\frac{1}{n(n+1)}(1+nx^n)^{1+\frac{1}{n}} + K$  (D)  $\frac{1}{n+1}(1+nx^n)^{1+\frac{1}{n}} + K$

Answer



53. The letters of the word **COCHIN** are permuted and all the permutations are arranged in an alphabetical order as in an English dictionary. The number of words that appear before the word **COCHIN** is
- (A) 360 (B) 192 (C) 96 (D) 48

Answer



54. Consider the planes  $3x - 6y - 2z = 15$  and  $2x + y - 2z = 5$ .


STATEMENT-1 : The parametric equations of the line of intersection of the given planes are  $x = 3 + 14t$ ,  $y = 1 + 2t$ ,  $z = 15t$ .

because

STATEMENT-2 : The vector  $14\hat{i} + 2\hat{j} + 15\hat{k}$  is parallel to the line of intersection of given planes.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer



55. STATEMENT-1 : The curve  $y = \frac{-x^2}{2} + x + 1$  is symmetric with respect to the line  $x = 1$ .

because

STATEMENT-2 : A parabola is symmetric about its axis.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer



(A)

(B)

(C)

(D)

56. Let  $f(x) = 2 + \cos x$  for all real  $x$ .

STATEMENT-1 : For each real  $t$ , there exists a point  $c$  in  $[t, t + \pi]$  such that  $f'(c) = 0$ .

because

STATEMENT-2 :  $f(t) = f(t + 2\pi)$  for each real  $t$ .

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer



(A)

(B)

(C)

(D)

57. Lines  $L_1 : y - x = 0$  and  $L_2 : 2x + y = 0$  intersect the line  $L_3 : y + 2 = 0$  at  $P$  and  $Q$ , respectively. The bisector of the acute angle between  $L_1$  and  $L_2$  intersects  $L_3$  at  $R$ .

STATEMENT-1 : The ratio  $PR : RQ$  equals  $2\sqrt{2} : \sqrt{5}$ .

because

STATEMENT-2 : In any triangle, bisector of an angle divides the triangle into two similar triangles.

- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
- (B) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
- (C) Statement-1 is True, Statement-2 is False
- (D) Statement-1 is False, Statement-2 is True

Answer ☐ ☐ ☒ ☐  
(A) (B) (C) (D)

58. Which one of the following statements is correct?

- (A)  $G_1 > G_2 > G_3 > \dots$
- (B)  $G_1 < G_2 < G_3 < \dots$
- (C)  $G_1 = G_2 = G_3 = \dots$
- (D)  $G_1 < G_3 < G_5 < \dots$  and  $G_2 > G_4 > G_6 > \dots$

Answer ☐ ☐ ☒ ☐  
(A) (B) (C) (D)

59. Which one of the following statements is correct?

- (A)  $A_1 > A_2 > A_3 > \dots$
- (B)  $A_1 < A_2 < A_3 < \dots$
- (C)  $A_1 > A_3 > A_5 > \dots$  and  $A_2 < A_4 < A_6 < \dots$
- (D)  $A_1 < A_3 < A_5 < \dots$  and  $A_2 > A_4 > A_6 > \dots$

Answer ☒ ☐ ☐ ☐  
(A) (B) (C) (D)

60. Which one of the following statements is correct?

(A)  $H_1 > H_2 > H_3 > \dots$

(B)  $H_1 < H_2 < H_3 < \dots$

(C)  $H_1 > H_3 > H_5 > \dots$  and  $H_2 < H_4 < H_6 < \dots$

(D)  $H_1 < H_3 < H_5 < \dots$  and  $H_2 > H_4 > H_6 > \dots$

**M61-63: Paragraph for Question Nos. 61 to 63**

If a continuous function  $f$  defined on the real line  $\mathbf{R}$ , assumes positive and negative values in  $\mathbf{R}$  then the equation  $f(x) = 0$  has a root in  $\mathbf{R}$ . For example, if it is known that a continuous function  $f$  on  $\mathbf{R}$  is positive at some point and its minimum value is negative then the equation  $f(x) = 0$  has a root in  $\mathbf{R}$ .

Consider  $f(x) = ke^x - x$  for all real  $x$  where  $k$  is a real constant.

Answer



(A)

(B)

(C)

(D)

61. The line  $y = x$  meets  $y = ke^x$  for  $k \leq 0$  at

(A) no point

(B) one point

(C) two points

(D) more than two points

Answer



(A)

(B)

(C)

(D)

62. The positive value of  $k$  for which  $ke^x - x = 0$  has only one root is

(A)  $\frac{1}{e}$

(B) 1

(C)  $e$

(D)  $\log_e 2$

Answer



(A)

(B)

(C)

(D)

63. For  $k > 0$ , the set of all values of  $k$  for which  $ke^x - x = 0$  has two distinct roots is

(A)  $\left(0, \frac{1}{e}\right)$

(B)  $\left(\frac{1}{e}, 1\right)$

(C)  $\left(\frac{1}{e}, \infty\right)$

(D)  $(0, 1)$

Answer



(A)

(B)

(C)

(D)

64. Let  $f(x) = \frac{x^2 - 6x + 5}{x^2 - 5x + 6}$ .

Match the expressions/statements in **Column I** with expressions/statements in **Column II** and indicate your answer by darkening the appropriate bubbles in the  $4 \times 4$  matrix given in the ORS.

**Column I**

**Column II**

(A) If  $-1 < x < 1$ , then  $f(x)$  satisfies

(p)  $0 < f(x) < 1$

(B) If  $1 < x < 2$ , then  $f(x)$  satisfies

(q)  $f(x) < 0$

(C) If  $3 < x < 5$ , then  $f(x)$  satisfies

(r)  $f(x) > 0$

(D) If  $x > 5$ , then  $f(x)$  satisfies

(s)  $f(x) < 1$

**Answer**

|   | p                                | q                                | r                                | s                                |
|---|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> |
| B | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            | <input checked="" type="radio"/> |
| C | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/>            | <input checked="" type="radio"/> |
| D | <input checked="" type="radio"/> | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> |

65. Let  $(x, y)$  be such that

$$\sin^{-1}(ax) + \cos^{-1}(y) + \cos^{-1}(bxy) = \frac{\pi}{2}.$$

Match the statements in **Column I** with statements in **Column II** and indicate your answer by darkening the appropriate bubbles in the  $4 \times 4$  matrix given in the ORS.

**Column I**

(A) If  $a = 1$  and  $b = 0$ , then  $(x, y)$

(B) If  $a = 1$  and  $b = 1$ , then  $(x, y)$

(C) If  $a = 1$  and  $b = 2$ , then  $(x, y)$

(D) If  $a = 2$  and  $b = 2$ , then  $(x, y)$

**Column II**

(p) lies on the circle  $x^2 + y^2 = 1$

(q) lies on  $(x^2 - 1)(y^2 - 1) = 0$

(r) lies on  $y = x$

(s) lies on  $(4x^2 - 1)(y^2 - 1) = 0$

**Answer**

|   | p                                | q                                | r                     | s                                |
|---|----------------------------------|----------------------------------|-----------------------|----------------------------------|
| A | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/> | <input type="radio"/>            |
| B | <input type="radio"/>            | <input checked="" type="radio"/> | <input type="radio"/> | <input type="radio"/>            |
| C | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/> | <input type="radio"/>            |
| D | <input type="radio"/>            | <input type="radio"/>            | <input type="radio"/> | <input checked="" type="radio"/> |

66. Match the statements in **Column I** with the properties in **Column II** and indicate your answer by darkening the appropriate bubbles in the  $4 \times 4$  matrix given in the ORS.

| Column I                                       | Column II                        |
|--|----------------------------------|
| (A) Two intersecting circles                   | (p) have a common tangent        |
| (B) Two mutually external circles              | (q) have a common normal         |
| (C) Two circles, one strictly inside the other | (r) do not have a common tangent |
| (D) Two branches of a hyperbola                | (s) do not have a common normal  |

Answer

|   | p                                | q                                | r                                | s                     |
|---|----------------------------------|----------------------------------|----------------------------------|-----------------------|
| A | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/> |
| B | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/>            | <input type="radio"/> |
| C | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |
| D | <input type="radio"/>            | <input checked="" type="radio"/> | <input checked="" type="radio"/> | <input type="radio"/> |